



Image 1 of 2

VIEW ENLARGEMENT



NO COPY

NDC 59267-1000-2

Pfizer-BioNTech COVID-19 Vaccine

Suspension for Intramuscular Injection

195 Multiple Dose Vials

(after dilution each vial contains 5 doses of 0.3 mL)



STORAGE: Prior to dilution, store at -80°C to -60°C (-112°F to -76°F). Store in this carton to protect from light.

DOSAGE AND ADMINISTRATION: After dilution, each vial contains 5 doses of 0.3 mL. See FDA-authorized Fact Sheet or scan QR code for information.

MUST BE DILUTED BEFORE USE with sterile 0.9% Sodium Chloride Injection, USP (not supplied). After dilution, store the vaccine at 2°C to 25°C (35°F to 77°F). Discard after 6 hours. Contains no preservative.

For use under Emergency Use Authorization.

Rx only



Manufactured by Pfizer Inc
New York, NY 10017
Manufactured for BioNTech
Manufacturing GmbH
An der Goldgrube 12
55131 Mainz, Germany



PAA156052

LOT:

EXP:

**IMPRINT AREA
(reads this way)
NO VARNISH
NO COPY, NO INK,**

NO COPY

NO COPY

Pfizer-BioNTech COVID-19 Vaccine

Suspension for Intramuscular Injection

195 Multiple Dose Vials

(after dilution each vial contains 5 doses of 0.3 mL)

STORAGE: Prior to dilution, store at -80°C to -60°C (-112°F to -76°F). Store in this carton to protect from light.

DOSAGE AND ADMINISTRATION: After dilution, each vial contains 5 doses of 0.3 mL. See FDA-authorized Fact Sheet or scan QR code for information.

MUST BE DILUTED BEFORE USE with sterile 0.9% Sodium Chloride Injection, USP (not supplied). After dilution, store the vaccine at 2°C to 25°C (35°F to 77°F). Discard after 6 hours. Contains no preservative.

PAA156052

LOT:

EXP:

**IMPRINT AREA
(reads this way)
NO VARNISH
NO COPY, NO INK,**

NO COPY

LOT/EXP

PAA156051

Pfizer-BioNTech COVID-19 Vaccine

After dilution, vial contains 5 doses of 0.3 mL

For intramuscular use. Contains no preservative.

For use under Emergency Use Authorization.

DILUTE BEFORE USE. Discard 6 hours after dilution when stored at 2 to 25°C (35 to 77°F).

Dilution date and time:

NDC 59267-1000-1

